



Root knot nematode management in pepper

Belay Feyisa, Fikremariam Yimer, Gemechu Kebede

Received: 16 December 2022 / Accepted: 18 May 2023 / Published: 30 June 2023

Ethiopian Institute of Agricultural Research Center (EIAR), Ambo Agricultural Research Center, P. O. Box 37, Ambo, Ethiopia.

*Correspondence

Belay Feyisa
belay22feyisa@gmail.com

Due to their slight negative effects on the environment, environmentally friendly nematode control techniques have been suggested. In order to determine the effectiveness of moringa leaf extracts and vermicompost alone and together against the root-knot nematodes, *Meloidogyne javanica* in pepper, the experiment was carried out in a greenhouse at the Ambo Agricultural Research Center., according to the results, 500 gm vermicompost and 80 gm moringa provided the greatest reduction in gall number per root (8), egg mass per 10 g root (2), gall index (2), population density per 100 g soil (108), and nematode reproduction factor (0.05). Vermicompost 500g and moringa 60g were also effective when combined. Additionally, organic amendments have a favorable effect on plant biomass. The findings obviously show that vermicomposting with plant products is more advantageous in organic farming. It is useful to control the damaging nematodes as well as for the improvement of soil nutrients.

Key words: *moringa, organic amendments, root-knot nematode, strategies, vermicompost*

Introduction

In Ethiopia, hot peppers (*Capsicum* spp.) are a common vegetable, spice, and condiment crop. The fruits are eaten fresh, dried, or processed. Among other vegetable crops, the production of peppers is a substantial source of foreign exchange and a strong source of revenue for small-scale producers or outside growers (Bonsu et al., 2003). It is a part of the typical diet of Ethiopians. 'Karia' (green pod), 'Berbere' (mature green fruits ground and mixed with other spices), and 'Mitmita' (the little, extremely pungent fruits) are significant ingredients in regional cuisine. According to estimates, an adult Ethiopian consumes 15g of hot pepper on a daily basis, more than tomatoes and the majority of other vegetables (MARC, 2004). Despite its significance, the output of hot peppers has remained modest, with national average yields of 6.2 t ha⁻¹ for green peppers and 1.7 t ha⁻¹ for dry pods (CSA, 2010). According to Eshetu et al. (2006), a variety of variables, including pests and a wide range of diseases, including root-knot nematodes, can reduce yield/unit of output and produce quality. RKNs are global treats and economically important PPNs of vegetable crops including pepper (Kokalis Burelle et al., 2009; Xing & Westphal, 2012). They cause extensive economic damage worldwide (Sikora & Fernandez2005). Published reports show that RKN (*Meloidogyne* spp.) is a serious and important pathogen of pepper which reduces the yield (Thies & Fery, 2002). Low levels of *Meloidogyne* in the soil can cause damage to different vegetables (pepper and tomato) and fruits (Wondiriad & Tesfamariam, 2002). The study undertaken by (Tadele & Mengistu, 2000) in Eastern Ethiopia revealed that infestation of RKN (*Meloidogyne* spp.) reaches up to 89% in fields. A national survey conducted by Ambo Agricultural Research Center revealed that the prevalence of RKN on pepper reaches 61.5% (unpublished). A number of methods were used for the management of RKN such as chemical control, organic amendments, resistant varieties, soil solarization and biological control (Agrios, 2005). RKN and other PPN can be effectively managed using nematicides (Nchore et al., 2011). Re-evaluation of these compounds has been prompted by worries about the damaging effects of synthetic nematicides on the environment and public health (Hassan et al., 2010; Udo & Ugwuoke, 2010). Alternative environmentally friendly approaches, such as organic amendments, have been suggested for the management of nematodes (Waceke, 2001 & 2002; Agyarko & Asante, 2005; Hassan et al., 2010; Nchore et al., 2011).

Phytochemicals of moringa have bio-nematicidal properties that affect the embryonic development of the nematode, inhibit, kill and or dissolve nematode egg mass (Adegbite, 2003, Agrio, 2005, D'Addabbo et al., 2010). Moringa leaf extracts reduce egg hatching of *M.incognita* up to (40-63.7%) and enhanced juvenile mortality from (82-93.8%) in the *invitro* (Claudius-Cole et al., 2010). So, identification of an effective, environmentally friendly, cost effective, easily available and applicable nematode management method is considered necessary. Therefore, this study aims to determine the effect of moringa and vermicompost on root-knot nematode (*M.javanica*).

Materials and methods

Inoculum preparation and treatment

Pure cultures of *M. javanica* raised from a single egg mass were used in the experiment. Infected roots were uprooted, rinsed free of soil, and cut into smaller pieces to allow the hatching of second stage juveniles for inoculation on new seedlings. For nematode reproduction, the most susceptible variety of tomato (cv. Melkasalsa) was used as the host plant to get enough inoculum for the experiment. Moringa leaves were collected from Wondogenet, shade dried, chopped and measured in grams (Sowley ENK., Kankam F., Adomako J., 2014). Vermicompost was obtained from Ambo Agricultural Research Center and measured in grams. Both treatments at different rates were used alone and in combination.

Table 1. Treatment arrangement and combinations

Treatment number	Treatment Type	Rate
T1	Vermicompost alone	500gm/pot
T2	Vermicompost alone	250gm/pot
T3	Moringa alone	80gm/pot
T4	Moringa alone	60gm/pot
T5	T1+T3	500+80gm/pot
T6	T1+T4	500+60gm/pot
T7	T2+T3	250+80gm/pot
T8	T2+T4	250+60gm/pot
Control	Only nematode	

Raising seedlings, nematode inoculation and treatment application

Fourteen days after treatment incorporation, four-week-old seedlings of a susceptible pepper cultivar (Oda hora) were transplanted on sterilized soil (2 kg per pot). One week after seedling transplantation, freshly hatched 2000-second juveniles (J2) of *Meloidogyne javanica* per pot were used for inoculation. The treatments were applied fifteen days before nematode inoculation. The control groups were inoculated without treatment. Pots were arranged in CRD with four replications. All agronomic practices were done uniformly for all pots. The experiment was terminated three months after inoculation of nematodes and plant growth parameters, including plant parameters, were recorded. The galls induced by *Meloidogyne javanica* were subsequently counted under a low power microscope (X6) and the degree of galling on a 0–5 scale (Taylor & Sasser 1978) was also determined at the end of the experiment. The nematode population in soil from each and every replicate was determined separately by processing 100 g of soil and 10 g of root using an extraction tray or a modified Baermann funnel method (Hooper et al., 2005).

Results and discussion

Effect of vermicompost and moringa leaf powder on growth parameters of pepper

The result revealed that there was a significant difference ($P < 0.05$) between the treatments on mean plant parameters (Table 1). Treatments also had an impact on plant shoots and root length and weight. The application of vermicompost significantly increased the length of shoots compared to the inoculated control treatment. This shows that the increment of pepper plant height by 28% over inoculated control. The application of vermicompost alone at 500gm/pot increases plant height by 25% over inoculated control. The highest plant height (116.80cm) was recorded in the combined application of vermicompost at (500gm) and Moringa at (80gm /pot) compared to control (91cm).

Plant height increased by 28 % over inoculated control treatments. The highest fresh shoot weight (316gm) was recorded from pots treated with combined application of vermicompost at (500gm) and moringa at (80gm /pot) compared to control (96.50gm). The highest fresh root weight (61gm) was recorded in pots treated with a combined application of vermicompost at (500gm) and moringa at (80gm/pot) compared to the control (14gm). Vermicomposting alone (250gm/pot) recorded the lowest fresh root weight compared to other treatments. The fresh root weight obtained from the control treatment was much less than that of other treatments including *Trichoderma harzianum* and aqueous moringa plant extracts. Female root-knot nematodes cause holes in roots, which cause the roots to lose water and moisture content. As a result, the weight of infected roots is lower than that of uninfected plants. This might have caused the fresh root weight of the control treatment to decrease. This finding was similar to Belay et al., (2019). Ahmad et al., 2007b and Siddiqui, 2006 were also reported that organic additives were effective in reducing RKN infestation and enhance various plant growth parameters and biomass. The inhibitory effect on nematodes combined with a direct fertilizing effect on the plants in the current study may explain why organic amendments improved plant development

Table 2. Effect of vermicompost and moringa leaf on height and biomass weight of pepper plants

Treatments	Shoot height(m)	Z**	Root length(m)	Z**	Fresh		Dry	
					Shoot weight(gm)	Root weight (gm)	Shoot weight (gm)	Root weight (gm)
Vermicompost alone (500gm/pot)	114.00 _{ab}	25	19.80 _b	88	262.50 _c	43.50 _c	55.80 _b	2.80 _c
Vermicompost alone(250gm/pot)	104.50 _d	15	12.5 _d	19	173.50 _f	38.00 _d	37.00 _e	2.20 _f
Moringa alone (80 gm/pot)	99.00 _e	9	12.00 _{de}	14	171.50 _f	33.00 _e	37.00 _e	1.20 _g
Moringa alone (60 gm/pot)	97.50 _e	7	11.30 _{ef}	8	161.50 _g	30.50 _f	35.50 _e	1.20 _g
Vermicompost 500 gm* Moringa 80 gm	116.8 _a	28	22.8 _a	117	318.00 _a	61.00 _a	61.3 _a	3.14 _a
Vermicompost 500 gm* Moringa 60 gm	115.00 _a	26	19.8 _b	88	274.30 _b	56.50 _b	56.8 _b	3.01 _b
Vermicompost 250gm* Moringa 80 gm	111.00 _{bc}	22	19. 5 _b	86	231.00 _d	42.50 _c	51.00 _c	2.6 _d
Vermicompost 250gm* Moringa 60 gm	109.00 _c	20	15.00 _c	43	215.00 _e	41.50 _c	46.80 _d	2.30 _e
Only nematode(2000J ₂)	91.00 _f		10.5 _f		96.50 _h	14.00 _g	25.00 _f	0.90 _h
CV	2.10		5.29		2.15	3.59	2.34	3.11
LSD_{0.05}	3.25		1.23		6.64	2.10	1.54	0.10

Values are averages of 4 replicates

Note: Means in column with the same letter are not significantly different (p< 0.05) DMRT.

Z** increase over inoculated control in percent

Effect of vermicompost and moringa leaf powder on root-knot nematode (*Meloidogyne javanica*)

The results showed that the treatments differed significantly (P 0.05) in terms of the mean number of galls per root, the number of eggs per root, the final nematode density per pot, and the reproduction factor (Table 2). In comparison to untreated control pots, pots treated with a combination of vermicompost (500gm) and moringa (80gm) had the lowest gall, egg mass, population density and reproduction factor. The pots treated with a combination of vermicomposting (500gm) and moringa (80gm) had the lowest population density (108) and reproduction factor

(0.05) as compared to the untreated control (3751, 1.90), respectively. It reduced the nematode population in soil by 97% and was significantly superior to other treatments. It was followed by Vermicompost (500gm) and Moringa (60gm), Vermicompost 250gm* Moringa 80 gm and only vermicompost 500gm. Vermicompost 250gm* Moringa 60 gm recorded the lowest reduction of soil nematode population, 27.2% over the inoculated control. The lower egg mass/root was also significantly less (2/10g root) in vermicomposting (500gm) and moringa (80gm) treatment as compared to the other treatments and inoculated control. Compared to the inoculated control, all the treatments minimize root gall development and reproduction of *M. javanica* on pepper plants. The reproduction factor ranged from 0.05 to 1.40 for the treated plants compared to 1.90 for the nematode control. Pots treated with vermicomposting (500gm) and moringa (80gm) showed 99% egg-mass reduction over inoculated control, which was in line with Claudius-Cole et al., 2010 reported that the Moringa leaf extracts reduce egg hatching of *M.incognita* up to (40-63.7%) and enhanced juvenile mortality from (82-93.8%) in *in-vitro*. According to the current study, plants that had their roots harmed by the nematode had a yellow color, were stunted, and were chlorotic (Figure 1, 2). Similar reports were made by Williamson & Gleason, 2003 and observable changes are made to the host's morphology and physiology by root-knot nematodes. Comparatively, all treatments have shown significant reduction in mean gall number and reproduction factor over the untreated control. Therefore, Vermicompost (500gm) & Moringa (80gm) and/or Vermicompost (500gm) and Moringa (60gm) are recommended for the management of Root Knot Nematodes; however, the study would better be repeated under field conditions for further confirmation and standardization.

Table 3. Effect of vermicompost and moringa leaf on root knot nematode (*Meloidogyne javanica*)

Treatments	No. of galls/root	No. of egg mass/root	Z**	Indices		Population density/pot	Z**	Reproduction factor
				Gall	Egg-mass			
Vermicompost alone (500gm/pot)	19.00 ^g	12.00 ^{fg}	95	2.80 _d	2.50 _d	364.00 ^f	90	0.18 ^f
Vermicompost alone(250gm/pot)	140.00 ^d	89.00 ^d	62	5.00 _a	4.00 _b	2137.00 ^c	76	1.07 ^c
Moringa alone (80 gm/pot)	180.00 ^c	130.00 ^c	44	5.00 _a	3.00 _b	2731.00 ^b	37	1.4 ^b
Moringa alone (60 gm/pot)	229.00 ^b	143.00 ^b	39	5.00 _a	5.00 _a	2732.00 ^b	37	1.4 ^b
Vermicompost 500 gm* Moringa 80 gm	8.00 ^h	2.00 ^g	99	2.00 _e	1.30 _f	108.00 ^h	97	0.05 ^h
Vermicompost 500 gm* Moringa 60 gm	14.00 ^{hg}	7.00 ^{gf}	97	3.00 _c	2.00 _e	294.00 ^g	92	0.15 ^g
Vermicompost 250gm* Moringa 80 gm	46.00 ^f	17.00 ^f	93	4.00 _b	3.00 _c	429.00 ^e	89	0.21 ^e
Vermicompost 250gm* Moringa 60 gm	121.00 ^e	61.00 ^e	74	5.00 _a	4.00 _b	1685.00 ^d	55	0.84 ^d
Only nematode(2000J ₂)	292.00 ^a	234.00 ^a		5.00 _a	5.00 _a	3751.00 ^a		1.90 ^a
LSD_{0.05}	9.86	11.14		0.24	0.36	50.10		0.01
CV	5.80	9.90		4.08	7.18	2.17		1.30

Values are averages of 4 replicates

Note: Means in column with the same letter are not significantly different (p< 0.05) DMRT.

Z** Reduction over inoculated control in percent



Figure 1. Variations b/n treatments on plant growth; Control (A), Vermicompost (500gm) *Moringa(80gm) (B) and Vermicompost(500gm) *Moringa(60gm) (C)



Figure 2. Variation between treatments on degree of root galling; T5=Vermicompost(500gm) *Moringa(80gm), T6=Vermicompost(500gm) *Moringa(60m), Ctl=Control

Conclusion

This study has demonstrated that the use of vermicomposting and moringa leaf against RKN has the potential for use as an economical, cost-effective method for managing RKNs in pepper production. While organic additives are safe for the environment, large-scale farming enterprises cannot use the technique due to the enormous quantities required per area. It is suggested that using plant residue in addition to other already employed management techniques might be more effective against nematodes. As a result of current research, it has been determined that the combination of vermicompost and *Moringa oleifera* proved to be quite successful in controlling *Meloidogyne javanica* in a greenhouse experiment. The treatment of two organic additives is more effective than individual organic additives.

Acknowledgement

The authors appreciate the financial and technical support provided by the Ethiopian Institutes of Agricultural Research, the Ambo Agricultural Research Center, Miss Alemnesh Zinaye and Miss Sinkinesh Jifara.

Author contributions

The research concept and the work proposal were created by Belay Feyisa. The experiment was carried out by Belay Feyisa, Fikeremariam Yimer, and Gemechu Kebede. They also analyzed the data and published the manuscript. The last draft was created by Belay Feyisa.

Competing interests

The authors have declared that no conflict of interest exists.

Ethics approval

Not applicable

References

- Adegbite, A.A. (2003). Comparative effects of carbofuran and water extract of *chromolonaodorata* on growth, yield and food component of root knot nematodes infested soybean (*Glycine max* L. Merrill) Ph.D. Diss., University of Ibadan, Ibadan, Nigeria. *J. Veg. Sci.*, 12, 5–12.
- Agrios, G.N. (2005). Introduction to Plant Pathology (5th Edition). Elsevier Academic Press Publication.
- Ahmad, F., Rather, M.A., & Siddiqui, M.A. (2007b). Impact of organic soil amendments and nematicides on *Meloidogyne incognita* infecting tomato. *Indian J. Nematol.*, 37, 55-57.
- Belay, F., Alemu, L., Thangavel, S., & Gezehegne, G. (2019). Effect of Some Botanicals and *Trichoderma Harzianum* against Root-Knot Nematode *Meloidogyne incognita*, Infecting Tomato under Green House. *Acad. Res. J. Agri. Sci. Res.* 7(5), 238-249.
- Bonsu, K.O., Owusu, E.O., Nkansah, G.O., Opong-Konadu, E., & Adu-Dapaah, H. (2003). Morphological characterization of hot pepper (*Capsicum* sp.) germplasm in Ghana. *Ghana J. Hortic.*, 2, 17-23.
- Claudius-Cole, A.O., Aminu, A.E., & Fawole, B. (2010). Evaluation of plant extracts in the management of root-knot nematode *Meloidogyne incognita* on cowpea [*Vigna unguiculata* (L) Walp]. *Mycopath.*, 8, 53-60.
- D'Addabbo, P., Avato, A., & Tava, E. (2010). Nematicidal potential of materials from *Medicago* spp.. *European Journal of Plant Pathology*, 125, 39-49.

- Esheteu, B., Ferdu, A., & Tsedeke, A. (2006). Experience with management of major plant diseases in Ethiopia pp. 25- 26. In Abrham T (eds) Proceedings on Facilitating the Implementation and Adoption of Integrated Pest Management (IPM) in Ethiopia. Melkassa Agricultural Research, October 13-15 2003, Ethiopian Agricultural Research Organization.
- Hassan, M., Chindo, P., & Alegbejo, M. (2010). Management of root knot nematodes (*Meloidogyne* spp.) on tomato (*Lycopersicon lycopersicum*) using organic wastes in Zaria, Nigeria. *Plant. Prot. Sci.*, 46,34-39.
- Hooper, D. U. (2005). Effects of biodiversity on ecosystem functioning: a consensus of current knowledge. *Ecol. Monogr.*, 75, 3 -35. <https://doi.org/10.1890/04-0922>
- Kokalis-Burelle, N., Bausher, M.G., & Roskopf, E.N. (2009). Green house evaluation of capsicum rootstocks for management of *Meloidogyne incognita* on grafted Bell pepper. *Nematropica*, 39, 121-132.
- MARC (Melkassa Agricultural Research Center). (2004). Progress Report, Addis Ababa, Ethiopia.
- Nchore, S.B., Waceke, W.J., & Kariuki, G.M. (2011). Incidence and Prevalence of root knot nematodes (*Meloidogyne species*) in selected indigenous leafy vegetables in Kisii and Transmara counties of Kenya. Department of plant and microbial sciences, Kenyatta University Kenya.
- Ogwulumba, S., Ugwuoke, K.I., & Ogbuji, R.O. (2010). Studies on *Meloidogyne javanica* infestation on roma tomato (*Lycopersicon esculentum* Mill.) under different soil amendment. *Afr. J. Biotec.*, 9, 3280-3283.
- Oostenbrink, M. (1966). Major characteristics of the relation between nematodes and plants. *Mededelingen*, 66,1-46.
- Siddiqui, M.A. (2006). Use of latex bearing plants to manage the nematodes attacking tomato and eggplant. *J.Eco-friendly Agric.*, 1,192-193.
- Sikora, R.A., & Fernandez, E. (2005). Nematode parasites of vegetables. Pp 319-392. In Luc, M., R. A. Sikora and J. Bridge (eds). *Plant-Parasitic Nematodes in Subtropical and Tropical Agriculture*. 2nd Edition. CAB International
- Sowley, E.N.K., Kankam, F., & Adomako J. (2014). Management of root-knot nematode (*Meloidogyne* spp.) on sweet pepper *Capsicum annum* L.) with moringa (*Moringa oleifera* Lam.) leaf powder. *Archives Phytopathol. Plant Prot.*, 47(13),1531-1538. <http://dx.doi.org/10.1080/03235408.2013.848710>
- Tadele, T., & Mengistu, H. (2000). Distribution of *Meloidogyne incognita* (Root knot nematode) in some vegetable's fields in eastern Ethiopia. *Pest Mgmt. J of Ethiopia*, 4,77-78.
- Taylor, A., & Sasser, J. (1978). *Biology, Identification and Control of Root-Knot Nematodes (Meloidogyne species)*. Department of Plant Pathology, North Carolina State University and U.S. Agency for International Development, Raleigh, NC, USA., 111p.
- Thies, J.A., & Fery, R.L. (2002). Evaluation of a core of the U.S. Capsicum germplasm for reaction to the northern root-knot nematode. *Hort. Science*, 37 (5), 805-810.
- Williamson, V.M., & Gleason, C.A. (2003). Plant nematode interactions. *Current Opinion in Plant Biology*, 6, 327-333.
- Wondirad, M., & Tesfamariam M. (2002). Root-knot nematodes on vegetable juveniles of *Meloidogyne incognita* in roots of *Arabidopsis thaliana*. *Nematology*, 38, 98-111.
- Xing, L., & Westphal, A. (2012). Predicting damage of *Meloidogyne incognita* on Watermelon. *Journal of Nematology*, 44 (2),127-133.